

Fragment-Based Restricted-Active-Space Configuration Interaction with Second-order Correction Embedding for Periodic Hartree-Fock Wave Function

Hung-Hsuan Lin¹, Denis Usvyat², Lorenzo Maschio³, Thomas Heine¹


¹Theoretical Chemistry, Technical University of Dresden, Germany.

²Institute for Chemistry, Humboldt University of Berlin, Germany.

³Department of Chemistry, University of Turin, Italy.

Keywords: excited states, electronic structure, configuration interaction, embedding scheme.

We present an embedding method for restricted-active-space configuration interaction with second-order perturbative correction (RASCI-PT2)¹ embedded in the periodic Hartree-Fock (HF) wave function. The method allows us to calculate the electronic structure of localized feature for fragment in periodic systems. We first outline the implementation of embedding scheme employing Crystal17², Cryscor³, and Q-Chem⁴. To evaluate our embedding method, we calculated the energy profile of single-bond dissociation of carbon-fluoride of fluorinated graphane, and the electronic excitation energies in covalent-organic framework. The results suggest that RASCI-PT2 embedded in HF wave function can be a promising tool for studying local properties of solid state.



1. Casanova, D. *J. Chem. Phys.* **2014**, 140.
2. R. Dovesi, A. Erba, R. Orlando, C. M. Zicovich-Wilson, B. Civalleri, L. Maschio, M. Rerat, S. Casassa, J. Baima, S. Salustro, B. Kirtman. *WIREs Comput Mol Sci.* **2018**, 8.
3. Pisani, C.; Schütz, M.; Casassa, S.; Usvyat, D.; Maschio, L.; Lorenz, M.; Erba, A. *Phys. Chem. Chem. Phys.* **2012**, 14.
4. Shao, Y.; Gan, Z.; Epifanovsky, E.; Gilbert, A. T. B.; Wormit, M.; Kussmann, J.; Lange, A. W.; Behn, A.; Deng, J.; Feng, X.; et al. *Mol. Phys.* **2015**, 113.